


# $\alpha$ clustering and flow in ultra-relativistic heavy-ion collisions

Wojciech Broniowski

Inst. of Nuclear Physics, Cracow, and  
Jan Kochanowski U., Kielce, Poland



SOTANCP3, Yokohama  
26-30 May 2014

[research with Enrique Ruiz Arriola, *arXiv:1312.0289*, PRL 112 (2014) 112501]

# Instead of outline

Two phenomena are related:

$\alpha$  clustering in light nuclei




harmonic flow in ultra-relativistic nuclear collisions

Surprising link:


lowest-energy ground-state structure  $\longleftrightarrow$  highest energy reactions

# $\alpha$ clusters

# $\alpha$ clusters expected in light nuclei



${}^9\text{Be}$



ground

Hoyle  $0^+$

other excited,  $2^+$  ...

${}^{12}\text{C}$

How can we detect the  $\alpha$  clusters in the ground state?



What is their spatial arrangement?

Assessment of  $n$ -body correlations (one-body not enough)

# Flow

# Ultra-relativistic A+A collisions (LHC, RHIC, SPS)


- Lorentz contraction
- Collision: essentially instantaneous passage, frozen configuration
- Reduction of the **ground-state** wave function of the nucleus (like measurement)



- detection of particles (mostly pions) in the transverse direction (mid-rapidity)

# Soft ( $p_T < 2$ GeV) particle production


Glauber-Czyż-Maximon model – particle production from independent NN collisions (wounded nucleon model of Białas, Błeszyński, and Czyż)



Au+Au collision at RHIC

# Soft ( $p_T < 2$ GeV) particle production

Glauber-Czyż-Maximon model – particle production from independent NN collisions (wounded nucleon model)



Au+Au collision at RHIC

- QCD partons produced in the initial phase  $\rightarrow$  sQGP


Subsequent phases:

- expansion
- freezeout
- hadrons (mostly pions) streaming to detectors





# Phenomenon of flow

Quark-gluon plasma is formed!



“Initial shape – final flow” transmutation detectable in the asymmetry of the momentum distribution of detected particles – follows from collectivity


# Elliptic flow



[ALICE]

$$dN/d\phi = A \left( 1 + 2 \sum_n v_n \cos[n(\phi - \Psi_n)] \right)$$


# Major observation in HIC – signature of QGP



[ALICE]

# Harmonic flow

Wounded nucleons –  
experienced at least one  
inelastic collision




- Initial fireball is asymmetric in the transverse plane from  
1) geometry 2) fluctuations
- **collectivity!** – **flow generated**
- Strong elliptic flow, **triangular** flow (in Au+Au entirely from fluctuations), higher-order harmonic flow

Merge the two ideas ( $\alpha$ 's and flow)  $\rightarrow$

# From $\alpha$ clusters to flow in relativistic collisions

$\alpha$  clusters  $\rightarrow$  asymmetry of shape  $\rightarrow$  asymmetry of initial fireball  $\rightarrow$   
 $\rightarrow$  hydro or transport  $\rightarrow$  collective harmonic flow

( $^{12}\text{C}$  as example)



nuclear triangular geometry  $\rightarrow$  fireball triangular geometry  $\rightarrow$  triangular flow


What are the signatures, chances of detection?  
(some blurring by fluctuations)

Related idea: triton/ $^3\text{He}$ -Au at RHIC in 2015

The case of  $^{12}\text{C}$  is more promising, as it leads to more abundant fireballs.

## Why ultrarelativistic?

Reaction time is much shorter than time scales of the structure  
→ a frozen “snapshot” of the nuclear configuration




wounding range determined by  $\sigma_{\text{NN}}^{\text{inel}}$

( $N_w > 70$  - flat-on orientation)

Imprints of the three  $\alpha$  clusters clearly visible


Simulations with GLISSANDO 2 – implementation of the **Glauber model**

... more events







... more events




... more events




... more events




... more events




... more events



... more events



... more events



## Intrinsic distributions

Ground state of  $^{12}\text{C}$  is a  $0^+$  state (rotationally symmetric wave function). The meaning of *deformation* concerns **multiparticle correlations** between the nucleons

Superposition over orientations:

$$|\Psi_{0^+}(x_1, \dots, x_N)\rangle = \frac{1}{4\pi} \int d\Omega \Psi_{\text{intr}}(x_1, \dots, x_N; \Omega)$$


The *intrinsic* density of sources of rank  $n$  is defined as the average over events, where the distributions in each event have aligned principal axes:  $f_n^{\text{intr}}(\vec{x}) = \langle f(R(-\Phi_n)\vec{x}) \rangle$ . Brackets indicate averaging over events and  $R(-\Phi_n)$  is the inverse rotation by the principal-axis angle in each event



# Our poor-man's $^{12}\text{C}$


[Could use actual distributions from theoretical calculations]

Three  $\alpha$ 's in a triangular arrangement, generate nucleon positions with Monte Carlo, parameters (size of the cluster, distance between clusters) chosen such that the EM form factor is reproduced




Electric charge density (dashed line) and the corresponding distribution of the centers of protons (solid line) in  $^{12}\text{C}$  for the data (i.e., after unfolding the proton charge form factor), plotted against the radius


Our intrinsic distributions in  $^{12}\text{C}$ : three  $\alpha$ 's in a triangular arrangement



*Intrinsic* distributions in the *transverse plane* of the fireball (here with  $N_w > 70$  – large multiplicity enforcing the flat-on collision)



clustered




unclustered

Some triangularity in the unclustered case follows from the fluctuations


# Geometry of nucleus $\rightarrow$ geometry of fireball

Triangular nucleus causes triangular "damage"!



intrinsic density of  $^{12}\text{C}$

$\rightarrow$



geometry of the fireball

# Eccentricity parameters

We need some quantitative measures of deformation (heavily used in heavy-ion analyses)

Eccentricity parameters  $\epsilon_n$  (Fourier analysis)

$$\epsilon_n e^{in\Phi_n} = \frac{\sum_j \rho_j^n e^{in\phi_j}}{\sum_j \rho_j^n}$$

describe the shape of each event ( $j$  labels the sources in the event,  $n$ =rank,  $\Phi_n$  is the principal axis angle)

$n = 2$  – ellipticity,  $n = 3$  – triangularity, ...

Two components:

- internal (from existent mean deformation of the fireball)
- from fluctuations

# Geometry vs multiplicity correlations in $^{12}\text{C}$ -Pb

Specific feature of the  $^{12}\text{C}$  collisions:

The cluster plane parallel or perpendicular to the transverse plane:




higher multiplicity  
higher triangularity  
lower ellipticity




lower multiplicity  
lower triangularity  
higher ellipticity

# Ellipticity and triangularity vs multiplicity



clustered



unclustered

Clusters: (qualitative signal)

When  $N_w \nearrow$  then  $\langle\epsilon_3\rangle \nearrow$  and  $\langle\epsilon_2\rangle \searrow$

and  $\langle\sigma(\epsilon_3)/\epsilon_3\rangle \searrow$ ,  $\langle\sigma(\epsilon_2)/\epsilon_2\rangle \nearrow$



No clusters:

similar behavior for  $n = 2$  and  $n = 3$

# Shape-flow transmutation

The eccentricity parameters are transformed (in all models based on collective dynamics) into asymmetry of the transverse-momentum flow  
It has been found that


$\langle v_n \rangle$  grows with  $\langle \epsilon_n \rangle$




→ for  $^{12}\text{C}$  collisions  $v_3$  will grow with multiplicity even stronger than  $\epsilon_3$   
[detailed modeling of  $v_3$  under way, research with Piotr Bożek]




# Dependence on the collision energy



$\sigma_{NN}^{inel} = 32\text{mb}$  (SPS)




42mb (RHIC)



72mb (LHC)






Qualitative conclusions hold from SPS to the LHC

# Other systems



# Other systems

(distributions matched to Wiringa's et al. radial densities)



[work with Maciej Rybczyński]



# Conclusions

# Nuclear structure from ultra-fast heavy ion collisions

**Snapshots** of the ground-state wave function

Spatial deformation of the ground state  $\rightarrow$  harmonic flow

**Signatures** (qualitative and quantitative) of **clustered**  $^{12}\text{C}$ - $^{208}\text{Pb}$  collisions

- Increase of triangularity with multiplicity for the highest multiplicity events
- Corresponding decrease of the scaled variance of triangularity
- Anticorrelation of ellipticity and triangularity

**Extensions**

- Other systems (e.g., in  $^{7,9}\text{Be}$  growth of  $v_2$  with multiplicity)
- More detailed modeling involving event-by-event hydrodynamics (in-progress) needed

Possible data (NA61@SPS, RHIC) in conjunction with a detailed knowledge of the evolution of the fireball would allow to place constraints on the  $\alpha$ -cluster structure of the colliding nuclei. Conversely, the knowledge of the clustered nuclear distributions helps to verify the fireball evolution