

# **Universality of hard-loop action**

**Stanisław Mrówczyński**

*Institute of Physics, Jan Kochanowski University, Kielce, Poland  
and National Centre for Nuclear Research, Warsaw, Poland*

in collaboration with **Alina Czajka**



# Outline

1. Motivation
2. Introduction
3. Keldysh-Schwinger formalism
4. Ghosts in Keldysh-Schwinger formalism
5. Self-energies
6. Universal hard-loop action
7. Limits of universality
8. Origin of universality
9. Summary

# Motivation

- 1997 – Maldacena – AdS/CFT duality (gauge/gravity correspondence)
- To what extent  $\mathcal{N}=4$  super YM is similar to QCD?
- To what extent  $\mathcal{N}=4$  super YM plasma is similar to quark-gluon plasma?
- To what extent a supersymmetric plasma is similar to its non-supersymmetric counterpart in a weak coupling regime?
  - Ph. D. thesis of Alina Czajka: [arXiv:1601.08215](https://arxiv.org/abs/1601.08215)

Two byproducts of the project:

- ghosts in the Keldysh-Schwinger formalism
- universality of the hard-loop action

# Gauge theories under consideration

$$\mathcal{L}_{\text{QED}} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + i \bar{\Psi} \gamma_\mu D^\mu \Psi$$

$$F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$$

$$\mathcal{L}_{\text{ScQED}} = -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} - (D_\mu \Phi)^* D^\mu \Phi$$

$$\begin{aligned} \mathcal{L}_{\text{SUSY QED}} = & -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + i \bar{\Psi} \gamma_\mu D^\mu \Psi + \frac{i}{2} \bar{\Lambda} \gamma_\mu \partial^\mu \Lambda + (D_\mu \phi_L)^* (D^\mu \phi_L) + (D_\mu^* \phi_R) (D^\mu \phi_R^*) \\ & + \sqrt{2} e (\bar{\Psi} P_R \Lambda \phi_L - \bar{\Psi} P_L \Lambda \phi_R^* + \phi_L^* \bar{\Lambda} P_L \Psi - \phi_R \bar{\Lambda} P_R \Psi) - \frac{e^2}{2} (\phi_L^* \phi_L - \phi_R^* \phi_R)^2 \end{aligned}$$

$$\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_a^{\mu\nu} F_{\mu\nu}^a$$

$$F_a^{\mu\nu} = \partial^\mu A_a^\nu - \partial^\nu A_a^\mu + g f^{abc} A_b^\mu A_c^\nu$$

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4} F_a^{\mu\nu} F_{\mu\nu}^a + i \bar{\Psi}_i (\gamma_\mu D^\mu)_{ij} \Psi_j$$

$$\begin{aligned} \mathcal{L}_{\text{SYM}} = & -\frac{1}{4} F_a^{\mu\nu} F_{\mu\nu}^a + \frac{i}{2} \bar{\Psi}_i^a (\gamma_\mu D^\mu \Psi_i)^a + \frac{1}{2} (D_\mu \Phi_A)_a (D^\mu \Phi_A)_a \\ & - \frac{1}{4} g^2 f^{abe} f^{cde} \Phi_A^a \Phi_B^b \Phi_A^c \Phi_B^d - i \frac{g}{2} f^{abc} (\bar{\Psi}_i^a \alpha_{ij}^p X_p^b \Psi_j^c + i \bar{\Psi}_i^a \beta_{ij}^p \gamma_5 Y_p^b \Psi_j^c) \end{aligned}$$

# Effective action

A system's dynamics is encoded in an effective action.

How to find the effective action?

Self-energy constrains a form of the effective action

$$\Pi^{\mu\nu}(x, y) = \frac{\delta^2 S[A]}{\delta A_\mu(x) \delta A_\nu(y)}$$

$$S = \int d^4x \mathcal{L}(x)$$

$$\mathcal{L}_2^{(A)}(x) = \frac{1}{2} \int d^4y A_\mu(x) \Pi^{\mu\nu}(x-y) A_\nu(y)$$

Our strategy



# Keldysh-Schwinger formalism

Description of non-equilibrium many-body systems


Contour Green function of gauge field

$$i\mathcal{D}_{ab}^{\mu\nu}(x, y) \stackrel{\text{def}}{=} \langle \tilde{T} A_a^\mu(x) A_b^\nu(y) \rangle$$

$$\langle \dots \rangle = \text{Tr}[\hat{\rho}(t_0) \dots]$$

$\tilde{T}$  - ordering along the contour


$$\tilde{T}A(x)B(y) = \Theta(x_0, y_0)A(x)B(y) \pm \Theta(y_0, x_0)B(y)A(x)$$




# Keldysh-Schwinger formalism

Contour Green's function includes 4 Green's functions  
with real time arguments:

$$\left(\mathcal{D}_{ab}^{\mu\nu}\right)^>(x, y) = \langle A_a^\mu(x) A_b^\nu(y) \rangle$$




$$\left(\mathcal{D}_{ab}^{\mu\nu}\right)^<(x, y) = \langle A_b^\nu(y) A_a^\mu(x) \rangle$$




$$\left(\mathcal{D}_{ab}^{\mu\nu}\right)^c(x, y) = \langle T^c A_a^\mu(x) A_b^\nu(y) \rangle$$

Chronological time ordering



$$\left(\mathcal{D}_{ab}^{\mu\nu}\right)^a(x, y) = \langle T^a A_a^\mu(x) A_b^\nu(y) \rangle$$

Anti-chronological time ordering



# Retarded, advanced & symmetric Green's functions

$$\mathcal{D}^+(x, y) = \Theta(x_0 - y_0) (\mathcal{D}^>(x, y) - \mathcal{D}^<(x, y))$$

$$\mathcal{D}^-(x, y) = \Theta(y_0 - x_0) (\mathcal{D}^<(x, y) - \mathcal{D}^>(x, y))$$

$$\mathcal{D}^{sym}(x, y) = \mathcal{D}^>(x, y) + \mathcal{D}^<(x, y)$$

# Meaning of the functions

$$\mathcal{D}^{<,>}(x, y)$$

- phase-space density
- mass-shell constraint
- real particles

$$\mathcal{D}^{\pm}(x, y)$$

- retarded & advanced propagator
- no mass-shell constraint
- virtual particles

The Green functions  $\mathcal{D}(x, y)$  are gauge dependent

**Physical results obtained from Green functions must be gauge independent**

For example

The poles of  $\mathcal{D}(x, y)$  - dispersion relations – are gauge independent

# Green's functions of free gluon field

$$D(x, y) = D(x - y)$$

Feynman gauge

$$\left(D_{\mu\nu}^{ab}\right)^>(p) = \frac{i\pi}{E_p} g_{\mu\nu} \delta^{ab} [\delta(E_p - p_0)[n_g(\mathbf{p}) + 1] + \delta(E_p + p_0)n_g(-\mathbf{p})]$$

$$\left(D_{\mu\nu}^{ab}\right)^<(p) = \frac{i\pi}{E_p} g_{\mu\nu} \delta^{ab} [\delta(E_p - p_0)n_g(\mathbf{p}) + \delta(E_p + p_0)[n_g(-\mathbf{p}) + 1]]$$

$$\left(D_{\mu\nu}^{ab}\right)^c(p) = -g_{\mu\nu} \delta^{ab} \left[ \frac{1}{p^2 + i0^+} - \frac{i\pi}{E_p} (\delta(E_p - p_0)n_g(\mathbf{p}) + \delta(E_p + p_0)n_g(-\mathbf{p})) \right]$$

$$\left(D_{\mu\nu}^{ab}\right)^a(p) = g_{\mu\nu} \delta^{ab} \left[ \frac{1}{p^2 - i0^+} + \frac{i\pi}{E_p} (\delta(E_p - p_0)n_g(\mathbf{p}) + \delta(E_p + p_0)n_g(-\mathbf{p})) \right]$$

$n_g(\mathbf{p})$  - gluon distribution function

# Green's functions of free gluon field

$$D(x, y) = D(x - y)$$

Feynman gauge

$$\left(D_{\mu\nu}^{ab}\right)^>(p) = \frac{i\pi}{E_p} g_{\mu\nu} \delta^{ab} [\delta(E_p - p_0)[n_g(\mathbf{p}) + 1] + \delta(E_p + p_0)n_g(-\mathbf{p})]$$

$$\left(D_{\mu\nu}^{ab}\right)^<(p) = \frac{i\pi}{E_p} g_{\mu\nu} \delta^{ab} [\delta(E_p - p_0)n_g(\mathbf{p}) + \delta(E_p + p_0)[n_g(-\mathbf{p}) + 1]]$$

$$\left(D_{\mu\nu}^{ab}\right)^+(p) = -\frac{g_{\mu\nu} \delta^{ab}}{p^2 + i \operatorname{sgn}(p_0) 0^+}$$

$$\left(D_{\mu\nu}^{ab}\right)^-(p) = \frac{g_{\mu\nu} \delta^{ab}}{p^2 - i \operatorname{sgn}(p_0) 0^+}$$

$n_g(\mathbf{p})$  - gluon distribution function

# Need for ghosts

- QCD computations in covariant gauges are usually much simpler than those in physical ones like the Coulomb gauge.
- Covariant gauges require ghosts to compensate unphysical degrees of freedom.

Gluon field:  $A^\mu(x) = (A^0(x), \mathbf{A}(x))$  - 4 degrees of freedom

Physical gluon: 2 polarizations

How to introduce ghosts in the Keldysh-Schwinger formalism?

What is the Green function of free ghosts?

$$\begin{array}{c} \Delta^>(p) \\ \Delta^<(p) \\ \Delta^c(p) \\ \Delta^a(p) \end{array} = ?$$

# How to get Green's function of free ghosts?

**Ghost sector should be determined by  
the gauge symmetry of the theory!**

$$A_\mu^a \rightarrow \left( A_\mu^a \right)^\psi = A_\mu^a + f^{abc} \omega^b A_\mu^c - \frac{1}{g} \partial_\mu \omega^a$$



gauge symmetry of the theory

**Slavnov-Taylor identities**

# Generating functional

$$W_0[J, \chi, \chi^*] = N_0 \int_{BC} \mathcal{D}A \mathcal{D}c \mathcal{D}c^* e^{i \int_C d^4x \mathcal{L}_{\text{eff}}(x)}$$

boundary conditions:  
the fields are fixed at  $t = -\infty \pm i0^+$

all fields are on the contour

$$\begin{aligned} \mathcal{L}_{\text{eff}}(x) = & -\frac{1}{4} F_a^{\mu\nu} F_{\mu\nu}^a - \frac{1}{2} (\partial^\mu A_\mu^a)^2 - c_a^* (\partial^\mu \partial_\mu \delta_{ab} - g \partial^\mu f^{abc} A_\mu^c) c_b \\ & + J_\mu^a A_\mu^\mu + \chi_a^* c_a + \chi_a c_a^* \end{aligned}$$

$$\begin{aligned} W[J, \chi, \chi^*] = & N \int DA' Dc' Dc^{*''} DA'' Dc'' Dc^{*''} \\ & \times \rho[A', c', c^{*''} | A'', c'', c^{*'''}] W_0[J, \chi, \chi^*] \end{aligned}$$

density matrix

# Generating functional

$$W[J, \chi, \chi^*] = N \int DA' Dc' Dc^{*''} DA'' Dc'' Dc^{*''} \\ \times \rho[A', c', c^{*''} | A'', c'', c^{*'''}] W_0[J, \chi, \chi^*]$$

The full Green's function are generated as

$$i\mathcal{D}_{\mu\nu}^{ab}(x, y) = (-i)^2 \frac{\delta^2}{\delta J_\mu^a(x) \delta J_\nu^b(y)} W[J, \chi, \chi^*] \Big|_{J=\chi=\chi^*=0}$$

The density matrix  $\rho[A', c', c^{*''} | A'', c'', c^{*'''}]$  is not specified



The explicit forms of the functional and the Green's function are not known

**The functional provides various relations among Green's functions**

# General Slavnov-Taylor identity

$$W[J, \chi, \chi^*] = N \int_{BC} \mathcal{D}A \Delta[A] e^{i \int_C d^4x \mathcal{L}(x)}$$



analog of the Fadeev-Popov determinant

$$\Delta[A] \equiv \int_{BC} \mathcal{D}c \mathcal{D}c^* e^{-i \int_C d^4x \left( -c_a^* (\partial^\mu \partial_\mu \delta_{ab} - g \partial^\mu f^{abc} A_\mu^c) c_b + \chi_a^* c_a + \chi_a c_a^* \right)}$$

The invariance of  $W[J, \chi, \chi^*]$  under the transformations

$$A_\mu^a \rightarrow \left( A_\mu^a \right)^U = A_\mu^a + f^{abc} \omega^b A_\mu^c - \frac{1}{g} \partial_\mu \omega^a$$

leads to

$$\left\{ i \partial_{(y)}^\mu \frac{\delta}{\delta J_d^\mu(y)} - \int_C d^4x J_a^\mu(x) \left( \partial_\mu^{(x)} \delta^{ab} + ig f^{abc} \frac{\delta}{\delta J_c^\mu(x)} \right) M_{bd}^{-1} \left[ \frac{1}{i} \frac{\delta}{\delta J} \Big| x, y \right] \right\} W[J, \chi, \chi^*] = 0$$

# Slavnov-Taylor identity for gluon Green's function

$$\frac{\delta}{\delta J_e^\nu(z)} \left\{ i\partial_{(y)}^\mu \frac{\delta}{\delta J_d^\mu(y)} - \int_C d^4x J_a^\mu(x) \left( \partial_\mu^{(x)} \delta^{ab} + ig f^{abc} \frac{\delta}{\delta J_c^\mu(x)} \right) M_{bd}^{-1} \left[ \frac{1}{i} \frac{\delta}{\delta J} \Big| x, y \right] \right\} W[J, \chi, \chi^*] = 0$$

$$J = \chi = \chi^* = 0$$

$$- p^\mu \mathcal{D}_{\mu\nu}^{ab}(p) = p_\nu \Delta_{ab}(-p)$$



free ghost Green's function

The longitudinal component of the gluon Green's function is free.

A. Czajka, St. Mrówczyński, PRD 89, 085035 (2014)

# Ghost functions

$$- p^\mu D_{\mu\nu}^{ab}(p) = p_\nu \Delta_{ab}(-p)$$

$$\Delta^>(p) = -\frac{i\pi}{E_p} \delta^{ab} \left[ \delta(E_p - p_0) [n_g(\mathbf{p}) + 1] + \delta(E_p + p_0) n_g(-\mathbf{p}) \right]$$

$$\Delta^<(p) = -\frac{i\pi}{E_p} \delta^{ab} \left[ \delta(E_p - p_0) n_g(\mathbf{p}) + \delta(E_p + p_0) [n_g(-\mathbf{p}) + 1] \right]$$


$$\Delta^c(p) = \delta^{ab} \left[ \frac{1}{p^2 + i0^+} - \frac{i\pi}{E_p} \left( \delta(E_p - p_0) n_g(\mathbf{p}) + \delta(E_p + p_0) n_g(-\mathbf{p}) \right) \right]$$

$$\Delta^a(p) = -\delta^{ab} \left[ \frac{1}{p^2 - i0^+} + \frac{i\pi}{E_p} \left( \delta(E_p - p_0) n_g(\mathbf{p}) + \delta(E_p + p_0) n_g(-\mathbf{p}) \right) \right]$$

$n_g(\mathbf{p})$  - gluon distribution function

# Polarization tensors

photon in QED



photon in scalar QED




photon in SUSY QED




gluon in Yang-Mills




gluon in QCD



gluon in  $\mathcal{N}=4$   
super Yang-Mills



# Contributions to gluon polarization tensor in QGP




$$\Pi_{ab}^{\mu\nu}(x, y) = -ig^2 N_c \delta_{ab} \text{Tr}[\gamma^\mu S(x, y) \gamma^\nu S(y, x)]$$


quark-loop contribution to contour polarization tensor

$S(x, y)$  - fermion contour Green's function


# Different loop contributions



$$\propto - \int d^3 p \frac{f(\mathbf{p})}{E_p} \left[ \frac{2p^\mu p^\nu + k^\mu p^\nu + p^\mu k^\nu - g^{\mu\nu} k \cdot p}{(p+k)^2 + i \text{sgn}((p+k)_0) 0^+} + \frac{2p^\mu p^\nu - k^\mu p^\nu - p^\mu k^\nu + g^{\mu\nu} k \cdot p}{(p-k)^2 - i \text{sgn}((p-k)_0) 0^+} \right]$$




$$\propto - \int d^3 p \frac{f(\mathbf{p})}{E_p} \left[ \frac{(2p+k)^\mu (2p+k)^\nu}{(p+k)^2 + i \text{sgn}((p+k)_0) 0^+} + \frac{(2p-k)^\mu (2p-k)^\nu}{(p-k)^2 - i \text{sgn}((p-k)_0) 0^+} \right]$$



$$\propto \int d^3 p \frac{f(\mathbf{p})}{E_p}$$

different structures  $\longrightarrow$  different behaviours

# Hard Loop Approximation



Wavelength of a quasi-particle is much bigger than inter-particle distance in the plasma:

$$\lambda \gg d$$

The only dimensional parameter in ultrarelativistic equilibrium plasma is temperature  $T$ , so:

$$\rho \sim \frac{1}{d^3} \sim T^3 \sim |\mathbf{p}|^3$$

$$p \sim \frac{1}{d} \quad k \sim \frac{1}{\lambda}$$

momentum at which a plasma is probed

$$k^\mu \ll p^\mu$$

momentum of plasma constituent

# Polarization tensor

$$\text{HL: } k^\mu \ll p^\mu$$

$$\Pi^{\mu\nu}(k) = C_\Pi \int \frac{d^3 p}{(2\pi)^3} \frac{f_\Pi(\mathbf{p})}{E_p} \frac{k^2 p^\mu p^\nu - [p^\mu k^\nu + k^\mu p^\nu - g^{\mu\nu}(k \cdot p)](k \cdot p)}{(k \cdot p + i0^+)^2}$$

After applying the HL approximation the polarization tensor gets **the same structure** for the  $\mathcal{N}=4$  SYM, YM, QCD, SUSY QED and usual QED plasma.

➤ **symmetric**

$$\Pi^{\mu\nu}(k) = \Pi^{\nu\mu}(k)$$






➤ **transversal**

$$k_\mu \Pi^{\mu\nu}(k) = 0$$

**Gauge independence!**

Plasma system	$C_\Pi$	$f_\Pi(\mathbf{p})$
QED	$e^2$	$2f_e(\mathbf{p}) + 2\bar{f}_e(\mathbf{p})$
Scalar QED	$e^2$	$f_s(\mathbf{p}) + \bar{f}_s(\mathbf{p})$
$\mathcal{N} = 1$ super QED	$e^2$	$2f_e(\mathbf{p}) + 2\bar{f}_e(\mathbf{p}) + 2f_s(\mathbf{p}) + 2\bar{f}_s(\mathbf{p})$
Yang-Mills	$g^2 N_c \delta^{ab}$	$2f_g(\mathbf{p})$
QCD	$g^2 N_c \delta^{ab}$	$2f_g(\mathbf{p}) + \frac{N_f}{N_c} (f_q(\mathbf{p}) + \bar{f}_q(\mathbf{p}))$
$\mathcal{N} = 4$ super Yang-Mills	$g^2 N_c \delta^{ab}$	$2f_g(\mathbf{p}) + 8f_f(\mathbf{p}) + 6f_s(\mathbf{p})$

# Fermionic self-energies

electron in QED	
electron in SUSY QED	 
photino in SUSY QED	
quark in QCD	
fermion in $\mathcal{N}=4$ super Yang-Mills	  

# Fermion self-energy


$$\Sigma(k) = C_\Sigma \int \frac{d^3 p}{(2\pi)^3} \frac{f_\Sigma(\mathbf{p})}{E_p} \frac{\hat{p}}{k \cdot p + i0^+}$$

HL:  $k^\mu \ll p^\mu$

The fermion self-energy in HL approximation has  
**the same structure** for all considered systems.

Plasma system	$C_\Sigma$	$f_\Sigma(\mathbf{p})$
QED	$\frac{e^2}{2}$	$2f_\gamma(\mathbf{p}) + f_e(\mathbf{p}) + \bar{f}_e(\mathbf{p})$
Electron in $\mathcal{N} = 1$ super QED	$\frac{e^2}{2}$	$2f_\gamma(\mathbf{p}) + f_e(\mathbf{p}) + \bar{f}_e(\mathbf{p}) + 2f_{\tilde{\gamma}}(\mathbf{p}) + f_s(\mathbf{p}) + \bar{f}_s(\mathbf{p})$
Photino in $\mathcal{N} = 1$ super QED	$\frac{e^2}{2}$	$f_e(\mathbf{p}) + \bar{f}_e(\mathbf{p}) + f_s(\mathbf{p}) + \bar{f}_s(\mathbf{p})$
QCD	$\frac{g^2 N_c^2 - 1}{2 \cdot 2N_c} \delta^{mn} \delta^{ij}$	$2f_g(\mathbf{p}) + N_f(f_q(\mathbf{p}) + \bar{f}_q(\mathbf{p}))$
$\mathcal{N} = 4$ super Yang-Mills	$\frac{g^2}{2} N_c \delta^{ab} \delta^{ij}$	$2f_g(\mathbf{p}) + 8f_f(\mathbf{p}) + 6f_s(\mathbf{p})$

# Scalar self-energies



$$P(k) = C_P \int \frac{d^3 p}{(2\pi)^3} \frac{f_P(\mathbf{p})}{E_p}$$

The scalar self-energy in HL approximation has **the same structure** for all considered systems.

Plasma system	$C_P$	$f_P(\mathbf{p})$
Scalar QED	$e^2$	$2f_\gamma(\mathbf{p}) + f_s(\mathbf{p}) + \bar{f}_s(\mathbf{p})$
$\mathcal{N} = 1$ super QED	$e^2$	$2f_\gamma(\mathbf{p}) + f_e(\mathbf{p}) + \bar{f}_e(\mathbf{p}) + 2f_{\tilde{\gamma}}(\mathbf{p}) + f_s(\mathbf{p}) + \bar{f}_s(\mathbf{p})$
$\mathcal{N} = 4$ super Yang-Mills	$g^2 N_c \delta^{ab} \delta^{AB}$	$2f_g(\mathbf{p}) + 8f_f(\mathbf{p}) + 6f_s(\mathbf{p})$

# From self-energies to effective action

The structure of self-energy of every field (vector, spinor, scalar) appears to be universal.

$$\Pi^{\mu\nu}(x, y) = \frac{\delta^2 S[A]}{\delta A_\mu(x) \delta A_\nu(y)} \quad S = \int d^4x \mathcal{L}(x)$$

$$\mathcal{L}_2^{(A)}(x) = C_\Pi \int \frac{d^3 p}{(2\pi)^3} \frac{f_\Pi(\mathbf{p})}{E_p} F_{\mu\nu}(x) \left( \frac{p^\nu p^\rho}{(p \cdot \partial)^2} \right) F_\rho^\mu(x)$$

$$\mathcal{L}_2^{(\Psi)}(x) = C_\Sigma \int \frac{d^3 p}{(2\pi)^3} \frac{f_\Sigma(\mathbf{p})}{E_p} \bar{\Psi}(x) \left( \frac{p \cdot \gamma}{p \cdot \partial} \right) \Psi(x)$$

$$\mathcal{L}_2^{(\Phi)}(x) = -C_P \int \frac{d^3 p}{(2\pi)^3} \frac{f_P(\mathbf{p})}{E_p} \Phi^*(x) \Phi(x)$$

$$\frac{1}{p \cdot \partial} \Psi(x) \equiv i \int \frac{d^4 k}{(2\pi)^4} \frac{e^{-ik \cdot x}}{p \cdot k} \Psi(k)$$

# Hard-loop effective action

$$\mathcal{L}_{\text{HL}}^{(A)}(x) = C_{\Pi} \int \frac{d^3 p}{(2\pi)^3} \frac{f_{\Pi}(\mathbf{p})}{E_p} F_{\mu\nu}(x) \left( \frac{p^\nu p^\rho}{(p \cdot D)^2} \right) F_{\rho}{}^{\mu}(x)$$

$$\mathcal{L}_{\text{HL}}^{(\Psi)}(x) = C_{\Sigma} \int \frac{d^3 p}{(2\pi)^3} \frac{f_{\Sigma}(\mathbf{p})}{E_p} \bar{\Psi}(x) \left( \frac{p \cdot \gamma}{p \cdot D} \right) \Psi(x)$$

$$\mathcal{L}_{\text{HL}}^{(\Phi)}(x) = -C_P \int \frac{d^3 p}{(2\pi)^3} \frac{f_P(\mathbf{p})}{E_p} \Phi^*(x) \Phi(x)$$

$$\frac{1}{p \cdot D} \Psi(x) \equiv \frac{1}{p \cdot \partial} \sum_{n=0}^{\infty} \left( i g p \cdot A(x) \frac{1}{p \cdot \partial} \right)^n \Psi(x)$$

E. Braaten, R. D. Pisarski, PRD 45, 1827 (1992)

St. Mrówczyński, A. Rebhan, M. Strickland, PRD 70, 025004 (2004)

**The structure of each term of the effective action appears to be unique.**

A. Czajka, St. Mrówczyński, PRD 91, 025013 (2015)

# Limitations of universality

## When is the universality valid?

Let us consider the limit  $k \rightarrow 0$

$$\Sigma(k) \sim g^2 \int \frac{d^3 p}{(2\pi)^3} \frac{f_\Sigma(\mathbf{p})}{E_p} \frac{\hat{p}}{k \cdot p + i0^+}$$

The wavevector  $k$  cannot be too small!

A. D. Linde, Phys. Lett. B 96, 289 (1980)

# The universality works when

# Physical consequences of universality

Microscopically different systems have the same long wavelength physical characteristics:

- response functions (dielectric function)
- screening lengths
- spectrum of collective modes (quasiparticles, instabilities)

## Dispersion equations

gauge boson field:  $\det[k^2 g^{\mu\nu} - k^\mu k^\nu - \Pi^{\mu\nu}(k)] = 0$

fermionic field:  $\det[\hat{k} - \Sigma(k)] = 0$

scalar field:  $k^2 + P(k) = 0$

# Why universality occurs?

Microscopic dynamics of different systems is **different**



Macroscopic behaviour of different systems is very **similar**

Simple example:

**QED plasma**

**vs.**

**scalar QED plasma**



Why is there no effect of **quantum statistics** of plasma constituents?

Why is there no effect of **different interactions**?

# Why universality occurs?

1. Hard loop condition:

$$k^\mu \ll p^\mu$$

$$\frac{1}{k} \text{ length scale at which the system is probed} \gg \frac{1}{p} \text{ de Broglie wavelength of plasma constituents}$$

**In classical limit fermions and bosons are not distinguishable!**

2. Gauge symmetry determines the interaction

# Summary

- The generating functional of QCD in the Keldysh-Schwinger formalism was constructed
- The general Slavnov-Taylor identity was derived
- The ghost Green's function was expressed through the gluon one
- QCD calculations in Keldysh-Schwinger approach are possible in the Feynman gauge
- Universal effective of various gauge theories was derived
- Universality results from long wavelength limit combined with gauge symmetry
- Properties of various plasma systems are very similar to each other