

Correlation of transverse momentum and multiplicity in a superposition model of nucleus-nucleus collisions

Stanisław Mrówczyński

*Institute of Physics, Świętokrzyska Academy, ul. Świętokrzyska 15, PL-25-406 Kielce, Poland
and Sołtan Institute for Nuclear Studies, ul. Hoża 69, PL-00-681 Warsaw, Poland*

(Received 14 February 2006; published 21 April 2006)

In p - p collisions the average transverse momentum is known to be correlated with the multiplicity of produced particles. The correlation is shown to survive in a superposition model of nucleus-nucleus collisions. When properly parametrized, the correlation strength appears to be independent of the collision centrality—it is the same in p - p and central A - A collisions. However, the correlation is strongly suppressed by the centrality fluctuations.

DOI: [10.1103/PhysRevC.73.044907](https://doi.org/10.1103/PhysRevC.73.044907)

PACS number(s): 25.75.Gz

I. INTRODUCTION

The transverse momentum of particles produced in p - p collisions is correlated with the particle's multiplicity. At the Super Proton Synchrotron (SPS) energy the correlation, which is directly observed experimentally [1,2], is most probably of simple kinematical origin—when the multiplicity N of produced particles at fixed collision energy grows, there is less and less energy to be distributed among transverse degrees of freedom of produced particles. Consequently, the average transverse momentum $\langle p_T \rangle_{(N)}$ decreases when N grows. The dependence of $\langle p_T \rangle_{(N)}$ on N is approximately linear [1,2].

The correlation p_T versus N seems to dominate the transverse momentum fluctuations, which have been experimentally studied [2] in nucleus-nucleus (A - A) collisions at SPS energy. The fluctuations have been quantified by the measure $\Phi(p_T)$ [3] and it has been shown [4] that the nonmonotonic behavior of $\Phi(p_T)$ as a function of collision centrality can be explained by the observed nonmonotonic multiplicity fluctuations [5,6] combined with the correlation p_T versus N . This contradicts the expectation that the correlation, which is well seen in p - p interactions, is washed out in A - A collisions. To clarify the situation the NA49 Collaboration has undertaken an effort to directly measure the correlation p_T versus N in A - A collisions at precisely determined numbers of projectile participants.

An analysis of the experimental data, which is underway, is performed for p - p and Pb-Pb collisions at 158A GeV in the forward rapidity window [4.0, 5.5] (with a center-of-mass rapidity of 2.92). The collision centrality is determined on an event-by-event basis, using a zero-degree calorimeter, which measures the energy deposited by remnants of the projectile nucleus. The energy is recalculated into the number of spectator nucleons coming from the projectile. The experimental method was worked out in the course of multiplicity fluctuation measurements performed at precisely fixed collision centralities [5,6].

The motivation of our analysis is twofold. The first goal, which is oriented toward experimental needs, is to check how a finite calorimeter resolution, which hinders the centrality determination, influences the observed correlation. The second goal is to formulate predictions within simple models of

nucleus-nucleus collisions as a reference for the experimental data. For these purposes we discuss the correlation p_T versus N in A - A collisions, assuming that the particles produced in such a collision originate from several identical sources that are independent of each other. A particular realization is the model where nucleus-nucleus collision is a superposition of independent nucleon-nucleon interactions. Then, a single nucleon-nucleon interaction is a single source of particles. More realistic is the wounded nucleon model [7] where the wounded nucleons—those that interact at least once—contribute independently to the final multiplicity. Then, there are two wounded nucleons in a nucleon-nucleon interaction, and thus, there are two particle sources. Since the particles in the forward domain are analyzed by the NA49 Collaboration, only the wounded nucleons from a projectile are expected to matter, and then their number equals the number of particle sources.

Our results obtained within the approach of independent sources can be interpreted by referring to one of three categories of models of nucleus-nucleus collisions introduced in Ref. [8]. One uses *transparency models* when the projectile (target) participants, which are excited in the course of interaction, mostly contribute to the particle production in the forward (backward) hemisphere. One deals with (rather unrealistic) *reflection models* when the projectile (target) participants mostly contribute to the backward (forward) hemisphere. Finally, there are *mixing models* when the projectile and target participants equally contribute to the backward and forward hemispheres. The latter models were shown [8,9] to properly describe (large) multiplicity fluctuations of particles produced in the forward hemisphere at fixed number of projectile participants [5,6], as the target participants of fluctuating number contribute to the forward hemisphere. The transparency models highly underestimate the multiplicity fluctuations whereas the reflection models strongly overestimate them.

Because of various possible interpretations of our results, we use in Secs. II–IV the term *source* but do not refer to a specific model. We first discuss the correlation p_T versus N for particles coming from a single source (Sec. II), and then, we consider particles that originate from several identical and independent sources (Sec. III). In Sec. IV a situation

with varying number of particle sources is discussed. Our results are interpreted in two models in Sec. V, where specific predictions are presented. To simplify the notation, the transverse momentum is denoted from now on as p , not as p_T or p_\perp . The word “transverse” is also mostly skipped in the text.

II. SINGLE SOURCE

$P_N(p_1, p_2, \dots, p_N)$ is the probability density that N particles with momenta p_1, p_2, \dots, p_N are produced by a single source. It is normalized as

$$\sum_N \int dp_1 dp_2 \cdots dp_N P_N(p_1, p_2, \dots, p_N) = 1. \quad (1)$$

The multiplicity distribution \mathcal{P}_N is given by

$$\mathcal{P}_N \stackrel{\text{def}}{=} \int dp_1 dp_2 \cdots dp_N P_N(p_1, p_2, \dots, p_N).$$

The momentum distribution of a single particle in the events of the multiplicity N is defined as

$$P_{(N)}(p) \stackrel{\text{def}}{=} \frac{1}{\mathcal{P}_N} \frac{1}{N} \left[\int dp_2 \cdots dp_N P_{(N)}(p, p_2, \dots, p_N) + \int dp_1 dp_3 \cdots dp_N P_{(N)}(p_1, p, p_3, \dots, p_N) + \cdots + \int dp_1 dp_2 \cdots dp_{N-1} P_{(N)}(p_1, p_2, \dots, p_{N-1}, p) \right], \quad (2)$$

and it is also normalized as

$$\int dp P_{(N)}(p) = 1.$$

The single-particle average momentum in the events with N particles, which is denoted as $\langle p \rangle_{(N)}$, is correlated with N , and this correlation has to be parametrized for our further considerations. According to the data [1,2], the dependence of $\langle p \rangle_{(N)}$ on N is approximately linear, and thus we write

$$\langle p \rangle_{(N)} = a + b \left(1 - \frac{N}{\langle N \rangle} \right). \quad (3)$$

We note that the parameter a only approximately equals the inclusive average momentum $\langle p \rangle$, which is

$$\langle p \rangle \stackrel{\text{def}}{=} \frac{1}{\langle N \rangle} \sum_N N \mathcal{P}_N \langle p \rangle_{(N)}, \quad (4)$$

with

$$\langle N \rangle \stackrel{\text{def}}{=} \sum_N N \mathcal{P}_N.$$

The parametrization (3) gives

$$\langle p \rangle = a + b \frac{\langle N \rangle^2 - \langle N^2 \rangle}{\langle N \rangle^2},$$

which for the case of poissonian multiplicity distribution is

$$\langle p \rangle = a - \frac{b}{\langle N \rangle}.$$

The multiplicity distribution of charged particles produced in high-energy p - p collisions is known not to be poissonian. First, the total number of charged particles is always even owing to charge conservation. The multiplicity distribution of positive or negative particles is not poissonian as well; the variance does not grow linearly with $\langle N \rangle$ but it follows the so-called Wróblewski formula [10] (i.e., the variance is a quadratic function of $\langle N \rangle$). However, for the collision energies of the SPS domain, which are of interest here, the Poisson distribution provides a reasonable approximation. The multiplicity distribution of not only positive or negative but of all charge particles is poissonian with high accuracy when the particles are registered in a limited acceptance. Therefore, the multiplicity distribution, which is further used in our calculation, is assumed to be of the Poisson form.

III. SEVERAL SOURCES

The momentum distribution of N particles coming from k identical and independent sources equals

$$P_N^k(p_1, p_2, \dots, p_M) = \sum_{N_1} \sum_{N_2} \cdots \sum_{N_k} \delta_N^{N_1+N_2+\cdots+N_k} \times P_{N_1}(p_1, p_2, \dots, p_{N_1}) P_{N_2}(p_{N_1+1}, p_{N_1+2}, \dots, p_{N_1+N_2}) \times \cdots \times P_{N_k}(p_{N_1+\cdots+N_{k-1}+1}, p_{N_1+\cdots+N_{k-1}+2}, \dots, p_{N_1+\cdots+N_k}). \quad (5)$$

One checks that the distribution (5) is normalized in agreement with Eq. (1). One defines the single-particle distribution analogous to (2) and computes the average single-particle momentum at fixed multiplicity as

$$\langle p \rangle_{(N)}^k = \frac{1}{\mathcal{P}_N^k} \frac{1}{N} \sum_{N_1} \sum_{N_2} \cdots \sum_{N_k} \delta_N^{N_1+N_2+\cdots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \cdots \mathcal{P}_{N_k} \times (N_1 \langle p \rangle_{(N_1)} + N_2 \langle p \rangle_{(N_2)} + \cdots + N_k \langle p \rangle_{(N_k)}). \quad (6)$$

Using the parametrization (3), one finds

$$\begin{aligned} \langle p \rangle_{(N)}^k &= \frac{1}{\mathcal{P}_N^k} \frac{1}{N} \sum_{N_1} \sum_{N_2} \cdots \sum_{N_k} \delta_N^{N_1+N_2+\cdots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \cdots \mathcal{P}_{N_k} \\ &\times \left[(N_1 + N_2 + \cdots + N_k)(a + b) - b \right. \\ &\times \left. \left(\frac{N_1^2}{\langle N_1 \rangle} + \frac{N_2^2}{\langle N_2 \rangle} + \cdots + \frac{N_k^2}{\langle N_k \rangle} \right) \right] \\ &= a + b - \frac{b}{\langle N \rangle} \frac{1}{\mathcal{P}_N^k} \frac{1}{N} \sum_{N_1} \sum_{N_2} \cdots \sum_{N_k} \delta_N^{N_1+N_2+\cdots+N_k} \\ &\times \mathcal{P}_{N_1} \mathcal{P}_{N_2} \cdots \mathcal{P}_{N_k} (N_1^2 + N_2^2 + \cdots + N_k^2), \quad (7) \end{aligned}$$

where it has been observed that

$$N_1 + N_2 + \cdots + N_k = N,$$

$$\langle N_1 \rangle + \langle N_2 \rangle + \cdots + \langle N_k \rangle = \langle N \rangle_k,$$

$$\mathcal{P}_N^k = \sum_{N_1} \sum_{N_2} \cdots \sum_{N_k} \delta_N^{N_1+N_2+\cdots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \cdots \mathcal{P}_{N_k}.$$

The average with the single-source distribution is denoted, as previously, as $\langle \dots \rangle$ whereas the average with the k -source distribution is $\langle \dots \rangle_k$.

Further calculation is performed with the poissonian multiplicity distribution, which is

$$\mathcal{P}_N = \frac{\langle N \rangle^N}{N!} e^{-\langle N \rangle}.$$

The convolution of k identical Poisson distributions of the average $\langle N \rangle$ is known to be equal to the Poisson distribution of the average $\langle N \rangle_k = k \langle N \rangle$. Thus,

$$\begin{aligned} & \sum_{N_1} \sum_{N_2} \dots \sum_{N_k} \delta_N^{N_1+N_2+\dots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \dots \mathcal{P}_{N_k} \\ &= \frac{(\langle N \rangle_k)^N}{N!} e^{-\langle N \rangle_k} = \mathcal{P}_N^k. \end{aligned} \quad (8)$$

Using the same technique, which provides Eq. (8), one proves that

$$\begin{aligned} & \sum_{N_1} \sum_{N_2} \dots \sum_{N_k} \delta_N^{N_1+N_2+\dots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \dots \mathcal{P}_{N_k} N_1^2 \\ &= \left[\frac{N}{k} + \frac{N(N-1)}{k^2} \right] \mathcal{P}_N^k. \end{aligned} \quad (9)$$

Substituting the formula (9) into Eq. (7), one finds the result

$$\langle p \rangle_{\langle N \rangle}^k = a + b \left(1 - \frac{N+k-1}{\langle N \rangle_k} \right), \quad (10)$$

which gives Eq. (3) for $k = 1$. If $\langle N \rangle \gg 1$, or equivalently $\langle N \rangle_k \gg k$, the explicit dependence of $\langle p \rangle_{\langle N \rangle}^k$ on k is effectively very weak. For the Poisson distribution with $\langle N \rangle_k \gg k \geq 1$, the multiplicities N of sizable probabilities do not differ much from $\langle N \rangle_k$. Consequently, $N+k-1$ can be approximated by N in Eq. (10) and we get the result analogous to the $k = 1$ case (3); that is,

$$\langle p \rangle_{\langle N \rangle}^k \approx a + b \left(1 - \frac{N}{\langle N \rangle_k} \right). \quad (11)$$

Thus, the average momentum at fixed multiplicity is approximately independent of the number of sources as it depends only on the ratio $N/\langle N \rangle_k$. The correlation strength parameter b is entirely independent of the source number.

IV. VARYING NUMBER OF SOURCES

In the previous section the number of particle sources has been treated as a fixed number. In a real experimental situation, however, the number of sources cannot be fully controlled in nucleus-nucleus collisions. The centrality selection is never perfect but even an exact determination of the participant number does not guarantee that the number of particle sources is always the same for a given number of participating nucleons. Thus, let us consider what happens when the number k varies event by event while the multiplicity of produced particles is kept constant.

When the number of particle sources fluctuates, the momentum distribution of N particles coming from k sources, which is given by Eq. (5), has to be averaged with the distribution of

number of sources \mathbf{P}_k . The analog of Eq. (6) reads

$$\begin{aligned} \langle \langle p \rangle_{\langle N \rangle}^k \rangle &= \frac{1}{\langle \mathcal{P}_N^k \rangle} \frac{1}{N} \sum_k \mathbf{P}_k \sum_{N_1} \sum_{N_2} \\ & \dots \sum_{N_k} \delta_N^{N_1+N_2+\dots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \dots \mathcal{P}_{N_k} \\ & \times (N_1 \langle p \rangle_{\langle N \rangle_{N_1}} + N_2 \langle p \rangle_{\langle N \rangle_{N_2}} + \dots + N_k \langle p \rangle_{\langle N \rangle_{N_k}}), \end{aligned} \quad (12)$$

where $\langle \mathcal{P}_N^k \rangle$ is the multiplicity distribution averaged over number of sources, which equals

$$\langle \mathcal{P}_N^k \rangle = \sum_k \mathbf{P}_k \sum_{N_1} \sum_{N_2} \dots \sum_{N_k} \delta_N^{N_1+N_2+\dots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \dots \mathcal{P}_{N_k}.$$

Substituting the parametrization (3) into Eq. (12), one finds

$$\begin{aligned} \langle \langle p \rangle_{\langle N \rangle}^k \rangle &= \frac{1}{\langle \mathcal{P}_N^k \rangle} \frac{1}{N} \sum_k \mathbf{P}_k \sum_{N_1} \sum_{N_2} \\ & \dots \sum_{N_k} \delta_N^{N_1+N_2+\dots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \dots \mathcal{P}_{N_k} \\ & \times \left[(N_1 + N_2 + \dots + N_k)(a + b) - b \right. \\ & \left. \times \left(\frac{N_1^2}{\langle N_1 \rangle} + \frac{N_2^2}{\langle N_2 \rangle} + \dots + \frac{N_k^2}{\langle N_k \rangle} \right) \right] \\ &= a + b - \frac{b}{\langle \mathcal{P}_N^k \rangle N \langle N \rangle} \sum_k \mathbf{P}_k k \sum_{N_1} \sum_{N_2} \\ & \dots \sum_{N_k} \delta_N^{N_1+N_2+\dots+N_k} \mathcal{P}_{N_1} \mathcal{P}_{N_2} \dots \mathcal{P}_{N_k} N_1^2. \end{aligned} \quad (13)$$

When the single-source multiplicity distribution \mathcal{P}_N is poissonian, one uses the formula (9), and Eq. (13) gives

$$\langle \langle p \rangle_{\langle N \rangle}^k \rangle = a + b \left(1 - \frac{1}{\langle N \rangle} - \frac{N-1}{\langle \mathcal{P}_N^k \rangle \langle N \rangle} \sum_k \mathbf{P}_k \frac{1}{k} \mathcal{P}_N^k \right). \quad (14)$$

Analytic calculations can be easily performed when the number of sources is treated not as an integer but as a continuous (positive) number distributed according to the gamma distribution

$$\mathbf{P}(k) = \frac{\alpha^{\beta+1}}{\Gamma(\beta+1)} k^\beta e^{-\alpha k}, \quad (15)$$

where $\Gamma(x)$ is the gamma function and the parameters α , β obey $\alpha > 0$ and $\beta > -1$. One easily finds the average and the variance of k as

$$\langle k \rangle = \frac{\beta+1}{\alpha}, \quad \sigma^2 = \frac{\beta+1}{\alpha^2}.$$

Substituting the Poisson multiplicity distribution (8) and the gamma distribution (15) into Eq. (14), one finds

$$\begin{aligned} \langle \langle p \rangle_{\langle N \rangle}^k \rangle &= a + b \left[1 - \frac{1}{\langle N \rangle} - \frac{\langle N \rangle \sigma^2 + \langle k \rangle}{\langle N \rangle} \right. \\ & \left. \times \frac{N-1}{(N-1)\sigma^2 + \langle k \rangle^2} \right]. \end{aligned} \quad (16)$$

Let us first discuss Eq. (16) in two limiting cases. When the variance of the source number σ^2 vanishes, Eq. (16) reproduces, as expected, the result (10) with $k = \langle k \rangle$. For $\sigma^2 \rightarrow \infty$, one gets

$$\langle \langle p \rangle_{\langle N \rangle}^k \rangle = a - \frac{b}{\langle N \rangle} = \langle p \rangle. \quad (17)$$

Thus, the momentum $\langle \langle p \rangle_{\langle N \rangle}^k \rangle$ is independent of N and it equals the inclusive average—the correlation of the transverse momentum and multiplicity entirely disappears.

The correlation of the transverse momentum and multiplicity is reduced owing to the source number fluctuation. To quantify the effect of the correlation reduction, Eq. (16) is expanded in powers of $(1 - N/\langle \langle N \rangle_k \rangle)$, where $\langle \langle N \rangle_k \rangle \equiv \langle k \rangle \langle N \rangle$, and only the linear term is kept. Then, Eq. (16) becomes

$$\langle \langle p \rangle_{\langle N \rangle}^k \rangle = a' + b' \left(1 - \frac{N}{\langle \langle N \rangle_k \rangle} \right), \quad (18)$$

where $a' \equiv a - 1/\langle N \rangle$ and

$$b' \equiv b \frac{\sigma^2 \langle N \rangle \langle k \rangle^3 + \langle k \rangle^4}{(\sigma^2 \langle N \rangle \langle k \rangle + \langle k \rangle^2 - \sigma^2)^2}. \quad (19)$$

When $\sigma^2 \rightarrow 0$, then $b' \rightarrow b$. For $\sigma^2 \rightarrow \infty$, $b' \rightarrow 0$. One also observes that $b' \rightarrow b$ for $\langle k \rangle \rightarrow \infty$. Equation (19) can be rewritten using the scaled variance $\omega \equiv \sigma^2/\langle k \rangle$ as

$$b' \equiv b \frac{\omega \langle N \rangle + 1}{(\omega \langle N \rangle + 1 - \omega/\langle k \rangle)^2},$$

but it does not eliminate an explicit dependence of b' on $\langle k \rangle$.

V. MODEL PREDICTIONS

We discuss here predictions of the formula (18), taking into account the conditions of the NA49 experiment where the particles produced in p - p and Pb-Pb collisions are observed, as mentioned in Sec. I, in the forward rapidity window [4.0, 5.5]. The coverage in the azimuthal angle is also incomplete. The average multiplicity of charged particles in the p - p interaction is $\langle N \rangle = 1.4$ in such an acceptance [2].

The formula (18) is first interpreted in the wounded nucleon model [7] belonging to the transparency class and, as explained in Sec. I, the source number k is identified with the number of wounded nucleons from a projectile N_{proj} . The number N_{proj} is experimentally determined on the event-by-event basis, using the zero-degree calorimeter. Owing to the finite energy resolution of the calorimeter, the number N_{proj} fluctuates with the dispersion identified as σ .

Figure 1 shows the ratio b'/b , which measures the correlation reduction, as a function of the average number of wounded nucleons $\langle N_{\text{proj}} \rangle$, for several values of the dispersion σ of the wounded nucleon number. As seen, the observable correlation strength is strongly reduced for the peripheral collisions and it does *not* match the p - p point ($b'/b = 1$ at $\langle N_{\text{proj}} \rangle = 1$), as one would naively expect.

The dispersion of N_{proj} is usually not constant but it changes with $\langle N_{\text{proj}} \rangle$. The energy resolution of the NA49 zero-degree calorimeter [11] that is recalculated into the dispersion of




FIG. 1. (Color online) The ratio b'/b as a function of the average number of projectile wounded nucleons $\langle N_{\text{proj}} \rangle$ in Pb-Pb collisions for several values of the dispersion σ of N_{proj} . The uppermost line corresponds to $\sigma = 1$, the lower one to $\sigma = 2$, etc.

projectile participant number N_{proj} is

$$\sigma_{\text{proj}}(\langle N_{\text{proj}} \rangle) = 0.77 \sqrt{A - \langle N_{\text{proj}} \rangle} - 0.28 - 0.032(A - \langle N_{\text{proj}} \rangle), \quad (20)$$

where A is the atomic number of the projectile nucleus. For the lead ($A = 208$) projectile, the formula (20) gives a σ that varies only between 3.5 and 4.0 for $1 < \langle N_{\text{proj}} \rangle < 150$, but it drops to 1.6 for $\langle N_{\text{proj}} \rangle = 200$. The prediction of the formula (19) with the dispersion of the wounded nucleon number given by Eq. (20) for $A = 208$ is represented by the uppermost curve labeled “res” in Fig. 2.

As mentioned in Sec. I, the transparency model, which assumes that the projectile participants mostly contribute to the particle production in the forward hemisphere, fails to describe [8,9] the large multiplicity fluctuations observed in the forward rapidity window [5,6]. The mixing models




FIG. 2. (Color online) b'/b vs $\langle N_{\text{proj}} \rangle$ in Pb-Pb collisions. The upper curve (“res”) shows the effect of calorimeter resolution; the middle one (“mix”) represents the mixing model; the lower curve (“res+mix”) combines the effect of calorimeter resolution and the mixing-model prediction.

[8,9], where the wounded nucleons from the projectile and target equally contribute to the forward rapidity window,¹ agree, at least qualitatively, with the data [5,6]. So, let us consider the model predictions concerning the correlation p_T versus N . The average number of sources $\langle k \rangle$ then equals $(\langle N_{\text{proj}} \rangle + \langle N_{\text{targ}} \rangle)/2$, where N_{targ} is the number of wounded nucleons from the target nucleus. Since we consider collisions of identical nuclei $\langle N_{\text{targ}} \rangle = \langle N_{\text{proj}} \rangle$, and thus $\langle k \rangle = \langle N_{\text{proj}} \rangle$.

We first neglect the experimental effect of finite calorimeter resolution, and we assume that N_{proj} is precisely determined; that is, N_{proj} does not fluctuate and $\langle N_{\text{proj}} \rangle = N_{\text{proj}}$. The number of particle sources, which contribute to the forward hemisphere, fluctuates owing to the fluctuations of N_{targ} at fixed N_{proj} and owing to the collision dynamics: A wounded nucleon from the projectile (target) is assumed to contribute with probability 1/2 to the forward (backward) hemisphere. Then, the variance of the particle sources, which contribute to the forward hemisphere, is $\sigma^2 = N_{\text{proj}}/2 + \sigma_{\text{targ}}^2/4$, where σ_{targ}^2 is the variance of N_{targ} at fixed N_{proj} . Using the numerical results obtained within the hadron-string dynamics model [12], which were used in Refs. [8,9], σ_{targ}^2 can be roughly parametrized as

$$\sigma_{\text{targ}}^2(N_{\text{proj}}) = 3.28 N_{\text{proj}} - 0.0158 N_{\text{proj}}^2$$

for the Pb-Pb collisions. As seen, σ_{targ} vanishes for $N_{\text{proj}} = 0$ and $N_{\text{proj}} = 208$, and it reaches a maximum equal to 13.1 at $N_{\text{proj}} = 104$. Thus, the variance of the particle sources is

$$\sigma^2(N_{\text{proj}}) = 1.32 N_{\text{proj}} - 0.004 N_{\text{proj}}^2, \quad (21)$$

with a maximum equal to 10.4 at $N_{\text{proj}} = 165$. Substituting $\langle k \rangle = N_{\text{proj}}$ and σ given by Eq. (21) into the formula (19), one finds the result shown in Fig. 2 by the middle curve labeled “mix.” As seen, the mixing model leads to a sizable reduction of the correlation p_T versus N , and the reduction does not depend much on the collision centrality.

Finally, we have combined the effect of “mixing” with that of the finite calorimeter resolution. Treating the two sources of fluctuations as independent of each other, the variances have been summed. The result is shown in Fig. 2 by the lowest curve labeled “res+mix.”

¹The analysis performed at RHIC energies shows that the contribution of a projectile (target) wound nucleon actually extends far into the backward (forward) hemisphere [13].

VI. SUMMARY AND CONCLUSIONS

The correlation p_T versus N , which is evident in p - p collisions, has been studied in the superposition model of nucleus-nucleus collisions where produced particles come from k identical and independent sources. At first the number of sources has been treated as a fixed number, and then the fluctuations of k have been taken into account. Although the superposition of particle sources preserves the correlation strength, the correlation is strongly reduced by the source number fluctuations. The calculations are fully analytical.

The wounded nucleon model has been used to formulate predictions for the NA49 measurement of the correlation p_T versus N in Pb-Pb collisions. Since the measurement, which is underway, is performed in the forward rapidity window, it has been first assumed that only wounded nucleons from the projectile contribute there. Taking into account a finite energy resolution of the zero-degree calorimeter, which allows one to precisely fix the number of wounded nucleons from the projectile, the correlation has been found to be strongly reduced in the peripheral collisions whereas the reduction in the central collisions is quite small.

Then, the “mixed” wounded nucleon model [8,9] has been considered. The model, which appears to be successful in describing the multiplicity fluctuations at a fixed number of projectile wounded nucleons [5,6], assumes, contrary to natural expectation [13], that the wounded nucleons from the projectile and target equally contribute to the forward rapidity window. In this case, the correlation p_T versus N seen in the forward rapidity window is strongly reduced and the reduction weakly depends on the collision centrality.

The correlation p_T versus N seems to be responsible [4] for the transverse momentum fluctuations [2] as quantified by the measure $\Phi(p_T)$ [3]. The data on the p_T versus N correlation combined with the existing data on the multiplicity [5,6] and transverse momentum fluctuations [2]—all obtained under the same experimental conditions—will hopefully allow one to formulate a coherent picture of the event-by-event fluctuations observed in nucleus-nucleus collisions.

ACKNOWLEDGMENTS

I am grateful to Marek Gaździcki for stimulating discussions and critical reading of the manuscript. Support by the Virtual Institute VI-146 of Helmholtz Gemeinschaft is also acknowledged.

-
- [1] T. Kafka *et al.*, Phys. Rev. D **16**, 1261 (1977).
 - [2] T. Antivicić *et al.* (NA49 Collaboration), Phys. Rev. C **70**, 034902 (2004).
 - [3] M. Gaździcki and St. Mrówczyński, Z. Phys. C **54**, 127 (1992).
 - [4] S. Mrówczyński, M. Rybczyński, and Z. Włodarczyk, Phys. Rev. C **70**, 054906 (2004).
 - [5] M. Gaździcki *et al.* (NA49 Collaboration), J. Phys. G **30**, S701 (2004).
 - [6] M. Rybczyński *et al.* (NA49 Collaboration), J. Phys. Conf. Ser. **5**, 74 (2005).
 - [7] A. Białaś, M. Błeszyński, and W. Czyż, Nucl. Phys. **B111**, 461 (1976).
 - [8] M. Gaździcki and M. Gorenstein, arXiv:hep-ph/0511058.
 - [9] V. P. Konchakovski, S. Haussler, M. I. Gorenstein, E. L. Bratkovskaya, M. Bleicher, and H. Stöcker, Phys. Rev. C **73**, 034902 (2006).
 - [10] A. Wróblewski, Acta Phys. Pol. B **4**, 857 (1973).
 - [11] M. Rybczyński, Ph.D. thesis, Kielce, 2005.
 - [12] W. Cassing and E. L. Bratkovskaya, Phys. Rep. **308**, 65 (1999).
 - [13] A. Białaś and W. Czyż, Acta Phys. Pol. B **36**, 905 (2005).