

# $\mathcal{N}=4$ super Yang-Mills Plasma

**Stanisław Mrówczyński**

*Institute of Physics,  
Jan Kochanowski University, Kielce, Poland  
and National Centre for Nuclear Research, Warsaw, Poland*

in collaboration with **Alina Czajka**




# Outline

1. Motivation
2.  $\mathcal{N}=4$  super Yang-Mills
3. Basic plasma characteristics
4. Collective modes
  1. Dispersion equations
  2. Perturbative computation of self-energies
  3. Effective action
5. Collisional characteristics
  1. Elementary processes
  2. Transport coefficients
  3. Energy loss &  $\hat{q}$
6. Conclusions

# Motivation


## AdS/CFT duality



QCD vs. super Yang-Mills?

## Motivation cont.

**Does rudimentary SUSY induce instabilities in fermionic sector?**



# Bigger project

A systematic comparison  
of supersymmetric plasma systems  
to their non-supersymmetric counterparts  
in a weak coupling domain

A. Czajka & St. Mrówczyński, Physical Review **D87** (2012) 065026  $\mathcal{N}=4$  super  
Yang-Mills

A. Czajka & St. Mrówczyński, Physical Review **D83** (2011) 065021 } SUSY  
A. Czajka & St. Mrówczyński, Physical Review **D84** (2011) 105020 } QED

# Lagrangian of $\mathcal{N}=4$ super Yang-Mills

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4} F_a^{\mu\nu} F_{\mu\nu}^a + \frac{i}{2} \bar{\Psi}_i^a (\mathcal{D}\Psi_i)^a + \frac{1}{2} (D_\mu \Phi_A)_a (D^\mu \Phi_A)_a \\ & - \frac{1}{4} g^2 f^{abe} f^{cde} \Phi_A^a \Phi_B^b \Phi_A^c \Phi_B^d \\ & - i \frac{g}{2} f^{abc} (\bar{\Psi}_i^a \alpha_{ij}^p X_p^b \Psi_j^c + i \bar{\Psi}_i^a \beta_{ij}^p \gamma_5 Y_p^b \Psi_j^c) \end{aligned}$$

Type of the field	Range of the field's index	Spin	Number of degrees of freedom
$A^\mu$ - vector	$\mu, \nu = 0, 1, 2, 3$	1	$2 \times (N_c^2 - 1)$
$\Phi_A$ - real (pseudo-)scalar	$A, B = 1, 2, \dots, 6$	0	$6 \times (N_c^2 - 1)$
$\lambda_i$ - Majorana spinor	$i, j = 1, 2, 3, 4$	$\frac{1}{2}$	$8 \times (N_c^2 - 1)$

# Basic plasma characteristics

	QGP	$\mathcal{N}=4$ SYMP
energy density - $\epsilon$	$\frac{\pi^2 T^4}{60} [4(N_c^2 - 1) + 7N_f N_c]$	$\frac{\pi^2 T^4}{2} (N_c^2 - 1)$
particle density - $n$	$\frac{2\zeta(3)T^3}{\pi^2} [2(N_c^2 - 1) + 3N_f N_c]$	$\frac{14\zeta(3)T^3}{\pi^2} (N_c^2 - 1)$
Debye mass - $m_D^2$	$\frac{g^2 T^2}{6} (2N_c + N_f)$	$2g^2 T^2 N_c$
plasma parameter - $\lambda$ $\left( \lambda \equiv \frac{1}{\frac{4}{3}\pi r_D^3 n} \right)$	$0.042 g^3$	$0.257 g^3$

All chemical potentials are assumed to vanish in both QGP and  $\mathcal{N}=4$  SYMP

# Collective modes



# Gluon dispersion equation

Equation of motion of gluon field  $A^\mu(k)$

$$[k^2 g^{\mu\nu} - k^\mu k^\nu - \Pi^{\mu\nu}(k)] A_\nu(k) = 0$$

$$k^\mu \equiv (\omega, \mathbf{k})$$

Dispersion equation

$$\det[k^2 g^{\mu\nu} - k^\mu k^\nu - \Pi^{\mu\nu}(k)] = 0$$

Collective modes - solutions:  $\omega(\mathbf{k})$

$\Pi^{\mu\nu}$  – retarded polarization tensor encodes gluon interaction with surrounding plasma

# Fermion & scalar dispersion equations

Fermion field

$$\det[k_{\mu}\gamma^{\mu} - \Sigma(k)] = 0$$

Scalar field

$$k^2 + P(k) = 0$$

# Keldysh-Schwinger formalism

Description of non-equilibrium many-body systems


Contour Green function of scalar field

$$iG(x, y) \stackrel{\text{def}}{=} \langle \tilde{T} \phi(x) \phi(y) \rangle$$

$$\langle \dots \rangle = \text{Tr}[\hat{\rho}(t) \dots]$$



$\tilde{T}$  - ordering along the contour

$$\tilde{T}A(x)B(y) = \Theta(x_0, y_0)A(x)B(y) \pm \Theta(y_0, x_0)B(y)A(x)$$





# Keldysh-Schwinger Green functions

Unordered functions – phase-space densities



Ordered functions – propagators



# Polarization tensor

Dyson-Schwinger equation


$$D = D_0 - D_0 \Pi D$$





$$D(x, y) = D(x - y) \text{ homogeneity, translational invariance}$$

# Lowest order contributions to $\Pi$

Contour-ordered Green functions have perturbative expansion similar to that of time-ordered Green functions




 gluon

 scalar

 fermion

 ghost

# Fermion-loop contribution to $\Pi$



Contour polarization tensor

$$\Pi_{ab}^{\mu\nu}(x, y) = -ig^2 N_c \delta_{ab} \text{Tr}[\gamma^\mu S(x, y) \gamma^\nu S(y, x)]$$

# From contour to retarded $\Pi$


$$\Pi^+(x, y) = \Theta(x_0 - y_0) (\Pi^>(x, y) - \Pi^<(x, y))$$

Contour polarization tensor

$$(\Pi(x, y))_{ab}^{\mu\nu} = -ig^2 N_c \delta_{ab} \text{Tr}[\gamma^\mu S(x, y) \gamma^\nu S(y, x)]$$

Unordered polarization tensor

$$(\Pi^>(x, y))_{ab}^{\mu\nu} = -ig^2 N_c \delta_{ab} \text{Tr}[\gamma^\mu S^>(x, y) \gamma^\nu S^<(y, x)]$$





# Fermion-loop contribution to $\Pi^+$

$$\begin{aligned}
 (\Pi^+(k))_{ab}^{\mu\nu} &= -\frac{ig^2}{2} N_c \delta_{ab} \int \frac{d^4 p}{(2\pi)^4} \times \\
 &\times \text{Tr}[\gamma^\mu S^+(p+k) \gamma^\nu S^{\text{sym}}(p) + \gamma^\mu S^{\text{sym}}(p) \gamma^\nu S^-(p-k)]
 \end{aligned}$$


Free Green functions

$$S^\pm(p) = \frac{p^\mu \gamma_\mu}{p^2 \pm ip_0 0^+} \quad S^{\text{sym}}(p) = S^>(p) + S^<(p)$$

$$S^>(p) = \frac{i\pi}{E_p} p^\mu \gamma_\mu [\delta(E_p - p_0)[n_f(\mathbf{p}) - 1] + \delta(E_p + p_0)n_f(-\mathbf{p})]$$


$$S^<(p) = \frac{i\pi}{E_p} p^\mu \gamma_\mu [\delta(E_p - p_0)n_f(\mathbf{p}) + \delta(E_p + p_0)[n_f(-\mathbf{p}) - 1]]$$

# Contributions to $\Pi$ in $\mathcal{N}=4$ SYMP



$$\left(\Pi^+(k)\right)_{ab}^{\mu\nu} = g^2 N_c \delta_{ab} \int \frac{d^4 p}{(2\pi)^4} \dots\dots$$

# Hard Loop Approximation



Wavelength of a quasi-particle is much bigger than inter-particle distance in the plasma

$$\lambda \gg d$$

## Hard Loop Approximation cont.

The only dimensional parameter in free ultrarelativistic equilibrium plasma is temperature  $T$ .

$$\text{particle density } \rho \sim \frac{1}{d^3} \sim T^3 \sim |\mathbf{p}|^3$$

$$\frac{1}{d} \sim |\mathbf{p}| \quad \text{momentum of plasma constituent}$$

$$\frac{1}{\lambda} \sim |\mathbf{k}| \quad \text{wave vector of collective mode}$$

**collective modes**

$$k^\mu \ll p^\mu$$

# HL polarization tensor

$$k^\mu \ll p^\mu$$

$$\Pi_{ab}^{\mu\nu}(k) = g^2 N_c \delta_{ab} \int \frac{d^3 p}{(2\pi)^3} \frac{f(\mathbf{p})}{E_p} \frac{k^2 p^\mu p^\nu - [p^\mu k^\nu + k^\mu p^\nu - g^{\mu\nu} (k \cdot p)] (k \cdot p)}{(k \cdot p + i0^+)^2}$$

$$f(\mathbf{p}) \equiv 2n_g(\mathbf{p}) + 8n_f(\mathbf{p}) + 6n_s(\mathbf{p})$$

➤ the same structure as in QED and QCD

➤ symmetric  $\Pi^{\mu\nu}(k) = \Pi^{\nu\mu}(k)$

➤ transversal

$$k_\mu \Pi^{\mu\nu}(k) = 0$$


**Gauge independence!**

## Effects of SUSY

- vacuum contribution vanishes ( $\Pi(k) = 0$  for  $f(\mathbf{p}) = 0$ )
- the coefficients in front of the distribution functions are the numbers of dof

$$f_{QGP}(\mathbf{p}) \equiv 2n_g(\mathbf{p}) + \frac{N_f}{N_c} (n_q(\mathbf{p}) + n_{\bar{q}}(\mathbf{p}))$$

# Fermion self-energy



$$\Sigma_{QED}(k) = e^2 \int \frac{d^3 p}{(2\pi)^3} \frac{f_\gamma(\mathbf{p}) + f_e(\mathbf{p})}{E_p} \frac{p^\mu \gamma_\mu}{k \cdot p + i0^+}$$

$$\Sigma_{ab}^{ij}(k) = \frac{g^2}{2} N_c \delta_{ab} \delta^{ij} \int \frac{d^3 p}{(2\pi)^3} \frac{f(\mathbf{p})}{E_p} \frac{p^\mu \gamma_\mu}{k \cdot p + i0^+}$$

$$f(\mathbf{p}) \equiv 2n_g(\mathbf{p}) + 8n_f(\mathbf{p}) + 6n_s(\mathbf{p})$$

The fermion self-energy has **the same structure** for the  $\mathcal{N}=4$  SYM, SUSY QED and usual QED plasma

# Scalar self-energy



$$P_{ab}^{AB}(k) = -2g^2 N_c \delta_{ab} \delta^{AB} \int \frac{d^3 p}{(2\pi)^3} \frac{f(\mathbf{p})}{E_p}$$

$$f(\mathbf{p}) \equiv 2n_g(\mathbf{p}) + 8n_f(\mathbf{p}) + 6n_s(\mathbf{p})$$

Scalar self-energy:

- independent of  $k$
- vanishes in the vacuum limit

# Hard loop effective action

Self-energy constrains the form of effective action

$$\mathcal{L}_2^{(\Psi)}(x) = \int d^4 y \bar{\Psi}(x) \Sigma(x-y) \Psi(y)$$

$$\Sigma(x, y) = \frac{\delta^2 S[\Psi, \bar{\Psi}]}{\delta \bar{\Psi}(x) \delta \Psi(y)}$$

$$\begin{aligned} \mathcal{L}_{\text{HL}} = & -\frac{1}{4} F_a^{\mu\nu} F_{\mu\nu}^a + \frac{i}{2} \bar{\Psi}_i^a (\not{D}\Psi_i)^a + \frac{1}{2} (D_\mu \Phi_A)_a (D^\mu \Phi_A)_a \\ & + \mathcal{L}_{\text{HL}}^{(A)} + \mathcal{L}_{\text{HL}}^{(\Psi)} + \mathcal{L}_{\text{HL}}^{(\Phi)} \end{aligned}$$



# From effective action to self-energies

$$\mathcal{L}_{\text{HL}}^{(A)}(x) = g^2 N_c \int \frac{d^3 p}{(2\pi)^3} \frac{f(\mathbf{p})}{E_p} F_{\mu\nu}^a(x) \left( \frac{p^\nu p^\rho}{(p \cdot D)^2} \right)_{ab} F_\rho^{b\mu}(x)$$

$$\mathcal{L}_{\text{HL}}^{(\Psi)}(x) = g^2 N_c \int \frac{d^3 p}{(2\pi)^3} \frac{f(\mathbf{p})}{E_p} \bar{\Psi}_i^a(x) \left( \frac{p \cdot \gamma}{p \cdot D} \right)_{ab} \Psi_i^b(x)$$

$$\mathcal{L}_{\text{HL}}^{(\Phi)}(x) = -2g^2 N_c \int \frac{d^3 p}{(2\pi)^3} \frac{f(\mathbf{p})}{E_p} \Phi_A^a(x) \Phi_A^a(x)$$

The structure of each term of the effective action appears to be unique

Structure of each self-energy is unique

# Gauge bosons collective modes

## Dispersion equation

$$\det[k^2 g^{\mu\nu} - k^\mu k^\nu - \Pi^{\mu\nu}(k)] = 0$$

solutions:  $\omega(\mathbf{k})$

$$k^\mu \equiv (\omega, \mathbf{k})$$

The structure of  $\Pi^{\mu\nu}(k)$

- coincides with the gluon polarization tensor of QCD plasma
- such as of QED and SUSY QED plasma

The spectrum of collective excitations of gauge bosons  
in  $\mathcal{N}=4$  super Yang-Mills, QCD, QED  
and SUSY QED plasma is the same

There is a whole variety of possible collective excitations,  
there are unstable modes

# Fermion collective modes

The structure of fermion self-energy is

- such as of quark self-energy in QCD plasma
- such as of QED and SUSY QED plasma

There are identical spectra of collective excitations  
of fermions in all systems

**No unstable modes found!**

Supersymmetry does not change anything

# Scalar collective modes

The self-energy is independent of momentum, negative and real

$$P(k) = -m_{\text{eff}}^2$$

$m_{\text{eff}}$  is the effective scalar mass

The solutions of dispersion equation

$$E_p = \pm \sqrt{m_{\text{eff}}^2 + \mathbf{p}^2}$$

# **Collisional characteristics**


# Elementary processes

Process	$\frac{1}{g^4} \frac{1}{N_c(N_c^2-1)} \sum  M ^2$
$GG \leftrightarrow GG$	$2\left(\frac{s^2+u^2}{t^2} + \frac{u^2+t^2}{s^2} + \frac{t^2+s^2}{u^2} + 3\right)$
$GF \leftrightarrow GF$	$32\left(\frac{s^2+u^2}{t^2} - \frac{u}{s} - \frac{s}{u}\right)$
$GG \leftrightarrow FF$	$32\left(\frac{t^2+u^2}{s^2} - \frac{u}{t} - \frac{t}{u}\right)$
$GS \leftrightarrow GS$	$24\left(\frac{s^2+u^2}{t^2} + 1\right)$
$GG \leftrightarrow SS$	$24\left(\frac{t^2+u^2}{s^2} + 1\right)$
$GF \leftrightarrow SF$	$-96\left(\frac{u}{s} + \frac{s}{u} + 1\right)$
$GS \leftrightarrow FF$	$-96\left(\frac{u}{t} + \frac{t}{u} + 1\right)$
$FS \leftrightarrow FS$	$-96\left[\frac{2us}{t^2} + 3\left(\frac{u}{s} + \frac{s}{u}\right) + 1\right]$
$SS \leftrightarrow FF$	$-96\left[\frac{2ut}{s^2} + 3\left(\frac{u}{t} + \frac{t}{u}\right) + 1\right]$
$SS \leftrightarrow SS$	$72\left(\frac{s^2+u^2}{t^2} + \frac{u^2+t^2}{s^2} + \frac{t^2+s^2}{u^2} + 3\right)$
$FF \leftrightarrow FF$	$128\left(\frac{s^2+u^2}{t^2} + \frac{u^2+t^2}{s^2} + \frac{t^2+s^2}{u^2} + 3\right)$

Coulomb-like scattering

# Transport coefficients

Collisional processes



**transport properties of ultrarelativistic plasma**

- ✓ Temperature  $T$  is the only dimensional parameter
- ✓ Coulomb-like scatterings dominate the interaction

shear viscosity  $\eta \sim \frac{T^3}{g^4 \ln g^{-1}}$

S. C. Huot, S. Jeon, and G. D. Moore, Phys. Rev. Lett. **98**, 172303 (2007)

# Energy loss & momentum broadening

- are not constrained by dimensional arguments

$$\frac{dE}{dx} \sim T^2, ET, E^2, \dots$$

$E$  – energy of test particle

$$\hat{q} \sim T^3, ET^2, E^2T, E^3, \dots$$

- depend on a specific scattering process under consideration




# Momentum broadening








Radiative energy loss of a fast parton is controlled by

$$\hat{q} \equiv \frac{d\langle \Delta p_T^2(t) \rangle}{dt}$$

Baier, Dokshitzer, Mueller, Peigne & Schiff 1996



# Elementary processes in SUSY QED

1	$e^{\mp}e^{\mp} \rightarrow e^{\mp}e^{\mp}$		$\frac{2\pi\alpha^2}{s^2} \left( \frac{s^2+u^2}{t^2} + \frac{s^2+t^2}{u^2} + \frac{2s^2}{tu} \right)$
2	$e^{\pm}e^{\mp} \rightarrow e^{\pm}e^{\mp}$		
3	$\gamma e^{\mp} \rightarrow \gamma e^{\mp}$		
4	$e^{\pm}e^{\mp} \rightarrow \gamma\gamma$		
5	$\gamma\gamma \rightarrow e^{\mp}e^{\pm}$		
23	$\tilde{e}_{L,R}^{\pm}\tilde{e}_{R,L}^{\mp} \rightarrow \tilde{e}_{R,L}^{\pm}\tilde{e}_{L,R}^{\mp}$		$\frac{\pi\alpha^2}{s^2}$
24	$\tilde{e}_{L,R}^{\mp}\tilde{e}_{L,R}^{\mp} \rightarrow \tilde{e}_{R,L}^{\mp}\tilde{e}_{R,L}^{\mp}$		$\frac{\pi\alpha^2}{s^2}$
25	$\gamma\tilde{e}_{L,R}^{\mp} \rightarrow \gamma\tilde{e}_{L,R}^{\mp}$		$\frac{4\pi\alpha^2}{s^2}$
26	$\tilde{e}_{L,R}^{\pm}\tilde{e}_{L,R}^{\mp} \rightarrow \gamma\gamma$		$\frac{8\pi\alpha^2}{s^2}$
27	$\gamma\gamma \rightarrow \tilde{e}_{L,R}^{\mp}\tilde{e}_{L,R}^{\pm}$		$\frac{2\pi\alpha^2}{s^2}$
31	$\tilde{e}_{L,R}^{\pm}e^{\mp}$		
32	$e^{\mp}e^{\mp}$		
33	$\tilde{e}_{L,R}^{\mp}\tilde{e}_{R,L}^{\mp} \rightarrow e^{\mp}e^{\mp}$		$\frac{4\pi\alpha^2}{s^2} \left( \frac{u}{t} + \frac{t}{u} \right)$

# Energy loss and momentum broadening in SUSY QED

A selectron is traversing an equilibrium photon gas.

$$|\mathcal{M}|^2 = 4e^4$$

$$\frac{dE}{dx} = -\frac{e^4}{2^5 3 \pi} T^2 \left[ 1 - \frac{12 \zeta(3) T}{\pi^2 E} \right] \underset{E \gg T}{\approx} -\frac{e^4}{2^5 3 \pi} T^2$$

$$\hat{q} = \frac{e^4}{12 \pi^3} T^3 \left[ \zeta(3) + \frac{\pi^4 T}{45 E} \right] \underset{E \gg T}{\approx} \frac{e^4 \zeta(3)}{12 \pi^3} T^3$$

# Comparison with Coulomb-like interaction


Energy loss for contact interaction

$$\frac{dE}{dx} = -\frac{e^4}{2^5 3\pi} T^2$$



Energy loss for Coulomb-like interaction

$$\frac{dE}{dx} = -\frac{e^4}{48\pi^3} T^2 \left( \ln \frac{E}{eT} + 2.031 \right)$$



E. Braaten and M. H. Thoma, Phys. Rev. D **44**, 1298 (1991)

# Energy loss

		Contact $ \mathcal{M} ^2 \sim e^4$	Coulomb $ \mathcal{M} ^2 \sim e^4 \frac{s^2}{t^2}$
energy change in single collision	$\Delta E$	$\sim E$	$\sim e^2 T$
cross section	$\sigma$	$\sim \frac{e^4}{ET}$	$\sim \frac{e^2}{T^2}$
density	$\rho$	$\sim T^3$	$\sim T^3$
inverse mean path	$\lambda^{-1} = \sigma\rho$	$\sim \frac{e^4 T^2}{E}$	$\sim e^2 T$
energy loss	$\frac{dE}{dx} \sim \frac{\Delta E}{\lambda}$	$\sim e^4 T^2$	$\sim e^4 T^2$

**Different interactions lead to the same energy loss!**

# Conclusions

- The collective modes of  $\mathcal{N}=4$  super Yang-Mills plasma are the same as those of QGP
  - The structures of self-energies appear to be unique
  - There are no unstable fermion modes
- The transport characteristics of SUSY plasma are similar to those of QGP

**Both systems are very similar to each other  
in the weak coupling regime!**